

ON THE BIMEROMORPHIC GEOMETRY OF COMPACT COMPLEX CONTACT THREEFOLDS

KRISTINA FRANTZEN AND THOMAS PETERNELL

ABSTRACT. We prove that a compact contact threefold which is bimeromorphically equivalent to a Kähler manifold and not rationally connected is the projectivised tangent bundle of a Kähler surface.

CONTENTS

1. Introduction	1
2. Uniruledness and splitting	2
3. The case of a 2-dimensional rational quotient	4
4. The case of a 1-dimensional rational quotient	8
References	11

1. INTRODUCTION

A (compact) complex manifold X of dimension $2n + 1$ is a *contact manifold* if there exists a vector bundle sequence

$$(\star) \quad 0 \rightarrow F \rightarrow T_X \xrightarrow{\theta} L \rightarrow 0,$$

where T_X is the tangent bundle and L a line bundle, with the additional property that the induced map $\bigwedge^2 F \rightarrow L$, $v \wedge w \mapsto [v, w]/F \in L$, given by the Lie bracket $[\cdot, \cdot]$ on T_X is everywhere non-degenerate. The line bundle L is referred to as the *contact line bundle* on X .

There are two basic ways to construct contact structures.

- A simple Lie group gives rise to a Fano contact manifold X (with $b_2(X) = 1$) by taking the unique closed orbit for the adjoint action of the Lie group on the projectivised Lie algebra, see e.g. [Bea98].
- For any compact complex manifold Y the projectivised tangent bundle $X = \mathbb{P}(T_Y)$ is a contact manifold.

Now the question naturally arises whether any compact complex contact manifold is given in this way.

In the following, let X be a compact complex contact manifold. If X is projective with $b_2(X) = 1$, then X must be a Fano manifold and Beauville [Bea98] proved

Date: May 10, 2010.

partial results towards the realisation as closed orbit. In general, if X is Kähler, Demailly [Dem02] showed that the canonical bundle K_X is not nef. If X is projective with $b_2(X) \geq 2$, Theorem 1.1 in [KPSW00] provides a positive answer to the question above. If X is Kähler but not projective, then necessarily $b_2(X) \geq 2$ and the second alternative is conjectured to hold, i.e., X should be a projectivised tangent bundle. However the paper [KPSW00] essentially uses Mori theory, which is, at the moment, not available in the Kähler case, except in dimension 3 where it can be shown that X is a projectivised tangent bundle over a surface (see Section 2).

In this paper we go one step further in dimension 3: we consider contact threefolds X which are in *class C*, i.e., bimeromorphic to a Kähler manifold. We first show that these threefolds must be uniruled. Then we consider the *rational quotient* $r : X \dashrightarrow Q$. The meromorphic map r identifies two very general points if and only if they can be joined by a chain of rational curves. In particular, X is *rationally connected* if and only if $\dim Q = 0$. We distinguish the cases $\dim Q = 1$ (Theorem 4.5) and $\dim Q = 2$ (Theorem 3.7) and show

Theorem. *Let X be a compact contact threefold in class $\mathcal{C}$. Assume that X is not rationally connected. Then there exists a Kähler surface Y such that $X \simeq \mathbb{P}(T_Y)$. In particular, X is Kähler.*

The remaining open case that X is rationally connected, in particular Moishezon, will require different methods. Probably it will be necessary to consider rational curves C with $-K_X \cdot C$ minimal, but positive.

2. UNIRULEDNESS AND SPLITTING

We shall use the following notation:

Definition 2.1. A compact complex manifold X is said to be in *class C* if X is bimeromorphically equivalent to a Kähler manifold.

An important property of manifolds in class $\mathcal{C}$ is the compactness of the irreducible components of the cycle space (cf. [Cam80])

The key for our investigations is the following

Theorem 2.2. *Let X be a compact contact threefold in class $\mathcal{C}$. Then X is uniruled.*

Proof. Let $\pi : \hat{X} \rightarrow X$ be a modification such that $\hat{X}$ is Kähler. It is a well-established fact that $\hat{X}$ is uniruled if and only if $K_{\hat{X}}$ is not pseudo-effective, i.e., the Chern class $c_1(K_{\hat{X}})$ is not represented by a positive closed current. The projective case in any dimension is treated in [BDPP04] based on the uniruledness theorem of Miyaoka-Mori [MM86]. The Kähler case in dimension three has been proved by Brunella [Bru06, Cor. 1.2].

The contact structure on X is given by $\theta \in H^0(X, \Omega_X^1 \otimes L)$; note that $\theta \wedge d\theta \neq 0$. Via π^* , the L -valued form θ induces a form $\hat{\theta} \in H^0(\hat{X}, \Omega_{\hat{X}}^1 \otimes \pi^*(L))$. By [Dem02, Cor. 1], the pullback of the dual line bundle $\pi^*(L^{-1})$ is not pseudo-effective. Since $K_X = 2L^{-1}$, the line bundle $\pi^*(K_X)$ is not pseudo-effective which is equivalent to say that K_X is not pseudo-effective. Since $\pi_*(K_{\hat{X}}) = K_X$, the Chern class

$c_1(K_{\hat{X}})$ cannot be represented by a positive closed current $\hat{T}$, because otherwise $c_1(K_X)$ would be represented by the positive closed current $\pi_*(\hat{T})$. Hence $K_{\hat{X}}$ is not pseudo-effective, and we conclude by the uniruledness criterion stated above. $\square$

As a consequence we obtain the following classification result for compact Kähler contact threefolds generalising the well-known projective case (see [KPSW00] for further references).

Corollary 2.3. *Let X be a compact Kähler contact threefold. Then either $X \simeq \mathbb{P}_3$ or $X = \mathbb{P}(T_Y)$ for a Kähler surface Y .*

Proof. By Theorem 2.2 above, the threefold X is uniruled. In particular, there is a positive-dimensional subvariety through the general point of X , i.e., X is not simple. The claim now follows from [Pet01, Theorem 4.1]. $\square$

Remark 2.4. The proof of Theorem 2.2 above actually shows that the canonical bundle of a compact contact manifold in class $\mathcal{C}$ of any dimension is not pseudo-effective. However in dimensions at least 4, unless X is projective, it is completely open, whether this implies uniruledness.

We now make a digression and consider the contact sequence $(\star)$. It is an interesting question whether this sequence can split. In the case where X is Fano, LeBrun [LeB95, Cor. 2.2] showed that splitting never occurs. By the following theorem, the same is true if X is in class $\mathcal{C}$.

Theorem 2.5. *Let X be a compact contact manifold in class $\mathcal{C}$. Then the contact sequence $(\star)$ does not split.*

Proof. Suppose we have a splitting $T_X \simeq F \oplus L$, hence $\Omega_X^1 \simeq F^* \oplus L^*$. Then it is well-known (see e.g. Beauville [Bea00]), that $c_1(L) \in H^1(X, L^*) \subset H^1(X, \Omega_X^1)$ and $c_1(F) \in H^1(X, F^*)$. Since $c_1(F) = nc_1(L)$ in $H^1(X, \Omega_X^1)$ and since X is in class $\mathcal{C}$, we conclude that $c_1(L) = c_1(F) = 0$ in $H^1(X, \Omega_X^1)$, and therefore also in $H^2(X, \mathbb{R})$. Hence K_X is numerically trivial and consequently, due to Remark 2.4, X cannot be in class $\mathcal{C}$. $\square$

Let X be a compact contact manifold X with contact sequence $(\star)$. A subvariety $S \subset X$, i.e., closed irreducible analytic subset in X , is called *F-integral* if $T_{S,x} \subset F_x$ for all smooth points $x \in S$.

Notation. A holomorphic family $(C_t)_{t \in T}$ of curves in X is given by a diagram

$$\begin{array}{ccc} Z & \xrightarrow{p} & X \\ \downarrow q & & \\ T & & \end{array}$$

such that

- T is an irreducible subspace of the cycle space of curves in X
- $q^{-1}(t)$ is the cycle corresponding to $t \in T$
- $C_t = p(q^{-1}(t))$ as cycle.

An important tool will be the following lemma, the proof of which is based on the observation that a surface covered by a family of F -integral curves is itself F -integral.

Lemma 2.6. *Let X be a compact contact threefold with contact line bundle L . Let $(C_t)_{t \in T}$ be a 1-dimensional family of generically irreducible rational curves passing through a fixed point $x_0 \in X$. Then $L \cdot C_t \geq 2$.*

Proof. We assume to the contrary that $L \cdot C_t \leq 1$ for all $t \in T$. By restricting the contact sequence $(\star)$ to an irreducible rational curve C_t and, if necessary, pulling it back to the normalisation $\eta : \tilde{C}_t \rightarrow C_t$, one finds that the map $\mathcal{O}(2) \simeq T_{\tilde{C}_t} \rightarrow \eta^*L \simeq \mathcal{O}(a)$ with $a \leq 1$ is trivial and therefore $T_{C_t, x} \hookrightarrow F_x$ for $x \in C_t$ smooth. I.e., for the general $t \in T$ the curve C_t is F -integral.

Consider the surface $S = \bigcup_{t \in T} C_t \subset X$ covered by the curves C_t . Then the proof of [Keb01, Proposition 4.1] shows that S is F -integral. But since any F -integral subvariety in X has dimension at most 1, this yields a contradiction. $\square$

3. THE CASE OF A 2-DIMENSIONAL RATIONAL QUOTIENT

We assume in this section that X is a compact contact threefold in class $\mathcal{C}$ with a rational quotient

$$r : X \dashrightarrow Q$$

of dimension $\dim Q = 2$. We refer the reader to the books [Deb01], [Kol96] and the references therein for relevant details on the construction and the properties of a rational quotient. A fundamental result of Graber, Harris, and Starr [GHS03] states that the quotient Q is not uniruled. In case X has dimension three, this result actually has previously been known.

The meromorphic map $r : X \dashrightarrow Q$ is almost holomorphic, i.e., r is proper holomorphic on a dense open set in X , and its general fiber is $\mathbb{P}_1$. Thus, we have a unique covering family $(l_t)_{t \in T}$ of rational curves with graph Z ,

$$\begin{array}{ccc} Z & \xrightarrow{p} & X \\ \downarrow q & & \downarrow r \\ T & & Q \end{array}$$

and $\dim T = 2$. Since r is almost holomorphic, the map p is bimeromorphic. By possibly passing to the normalisation, we may assume both Z and T normal. Moreover, we may take $Q = T$.

Lemma 3.1. *All curves l_t satisfy $L \cdot l_t = 1$ and all irreducible curves l_t are F -integral.*

Proof. The general l_t is a general fiber of r , hence $-K_X \cdot l_t = 2$ by adjunction and therefore $L \cdot l_t = 1$. The same is then true for all l_t . All irreducible curves l_t are consequently F -integral (cf. proof of Lemma 2.6). $\square$

In the following we will make use of the deformation theory of rational curves. This is to say we consider a rational curve $C \subset X$, given by a bimeromorphic morphism $f : \mathbb{P}_1 \rightarrow X$ and consider the deformations f_t of f . We obtain a family $(C_t) = (f_t(\mathbb{P}_1))$ and then take its closure in the cycle space, because in general the family (f_t) will not be compact, or in other words, the family (C_t) will split. Here it is essential that X is in class $\mathcal{C}$, hence all irreducible components of the cycle space of X are compact. We repeatedly use the following basic fact, see e.g. [Kol96, Theorem II.1.3].

Fact 3.2. *Let X be a compact threefold and let C be a rational curve in X . If $-K_X \cdot C \geq m$, then C will deform as rational curve in an at least m -dimensional family.*

The following proposition is the technical core of this section.

Proposition 3.3. *The map $p : Z \rightarrow X$ is an isomorphism. In particular, the rational quotient $r : X \rightarrow T$ is holomorphic and equidimensional.*

Proof. For $x \in X$ we let $T(x)$ be the analytic subset of all $t \in T$ such that $x \in l_t$. Since the general l_t does not pass through x , it follows that $\dim T(x) \leq 1$. In the following we show that $\dim T(x) = 0$ for all x ; in other words, p is finite. Since the map p is of degree 1 and has connected fibers by Zariski's main theorem, the finiteness of p forces p to be biholomorphic.

Suppose now to the contrary that $\dim T(x) = 1$ for some fixed $x \in X$. If $T(x)$ happens to be reducible, we replace it by an irreducible component of dimension 1. In the following, we shall therefore assume that $T(x)$ is irreducible.

Let S be the surface covered by the l_t belonging to $T(x)$:

$$S = \bigcup_{t \in T(x)} l_t.$$

If the general l_t through x is irreducible, then the Lemmata 2.6 and 3.1 yield a contradiction.

So we are left with the case when all l_t , $t \in T(x)$ are reducible. In this case S itself might be reducible. For $t \in T(x)$ we decompose l_t into its irreducible components and write $l_t = \sum a_t^j C_t^j$. Since $L \cdot l_t = 1$ for all $t \in T$, there exists at least one component C_t^j in this decomposition with $L \cdot C_t^j \geq 1$. We pick $t \in T(x)$ general and let $C^{(1)}$ be a component of l_t with $L \cdot C^{(1)} \geq 1$. Then by Fact 3.2, $C^{(1)}$ deforms in an at least 2-dimensional family $(C_t^{(1)})_{t \in T_1}$.

Suppose first that

$$L \cdot C_t^{(1)} = 1.$$

If the family $(C_t^{(1)})_{t \in T_1}$ covers a surface, then we find a 1-dimensional subfamily through a fixed point, contradicting Lemma 2.6. If the family covers all of X , then, since there is only one covering family of generically irreducible rational curves in X , the family $(C_t^{(1)})_{t \in T_1}$ must be the original family $(l_t)_{t \in T}$, in particular $T = T_1$. In other words, we have $t_0 \in T(x)$ and $t_1 \in T$ such that $l_{t_0} = l_{t_1} + R$ with an effective curve R . Thus $p^{-1}(x)$ contains more than one point for every $x \in l_{t_1}$. Since p has connected fibers, we conclude that $\dim p^{-1}(x) = 1$ for every $x \in l_{t_1}$. Then either all curves l_t , $t \in T$ pass through l_{t_1} , which is absurd since r is almost

holomorphic, or there exists a 1-dimensional subfamily $(l_{t_1} + C_u)_{u \in U}$ of $(l_t)_{t \in T}$ with $\dim U = 1$. In this second case however, since the subfamily $(l_{t_1} + C_u)_{u \in U}$ does not contain the curve l_{t_1} itself, it follows $p^{-1}(x)$ is not connected, a contradiction.

So we are left with

$$L \cdot C_t^{(1)} \geq 2,$$

i.e., $-K_X \cdot C_t^{(1)} \geq 4$, and the $C_t^{(1)}$ deform in an at least 4-dimensional family (cf. Fact 3.2) which must stay in an irreducible component S_1 of S . (The deformations of $C_t^{(1)}$ cannot cover all of X since there is a unique covering family of generically irreducible rational curves in X , and this family, namely $(l_t)_{t \in T}$, is 2-dimensional and fulfils $-K_X \cdot l_t = 2$). We want to exhibit a new family $(C_t^{(2)})$ in the surface S_1 such that $L \cdot C_t^{(2)} \geq 2$.

In order to construct this new family we notice that the 4-dimensional family $(C_t^{(1)})_{t \in T_1}$ must split. In fact, through any two points of S_1 there is a positive-dimensional subfamily. Now we choose carefully a splitting component $C^{(2)}$ such that $L \cdot C^{(2)} \geq 2$, namely we want to achieve that $C^{(2)}$ passes through a general point of S_1 . By Lemma 3.5 we obtain a generically non-splitting family $(h_u)_{u \in U}$ of rational curves h_u in S_1 with $\dim U \geq 2$ such that for general $u \in U$ there exists $t(u) \in T_1$ such that h_u is an irreducible component of $C_{t(u)}^{(1)}$. Since the family (h_u) covers exactly S_1 , there exists a 1-dimensional subfamily through a general point of S_1 and we take $C^{(2)}$ to be a general member of this subfamily. By Lemma 2.6 we obtain

$$L \cdot C^{(2)} \geq 2.$$

Again, $-K_X \cdot C^{(2)} \geq 4$ implies that $C^{(2)}$ moves in an at least 4-dimensional family of rational curves, say $(C_t^{(2)})_{t \in T_2}$.

Inductively we obtain families $(C_t^{(k)})_{t \in T_k}$ in S_1 such that

$$L \cdot C_t^{(k)} \geq 2.$$

Our aim now is to find an argument that this procedure must stop at some point, i.e., that $L \cdot C_t^{(k)} \geq 2$ cannot occur infinitely many times.

If X is Kähler with Kähler form ω , this follows from the fact that the intersection number $C_t^{(k)} \cdot \omega$ strictly decreases and that all classes $C_t^{(k)}$ are *integer* classes in a ball $\{a \mid a \cdot \omega \leq K\} \subset H^2(X, \mathbb{R})$.

Let us briefly explain the difficulty arising from the fact that X is not necessarily Kähler. If X is merely in class $\mathcal{C}$, we cannot argue in this way, because we will have curves with “semi-negative” cohomology. To be precise, we choose a sequence of blow-ups in points and smooth curves $\pi : \hat{X} \rightarrow X$ such that $\hat{X}$ is Kähler, fix a Kähler form $\hat{\omega}$ on $\hat{X}$, and form the current $R = \pi_*(\omega)$. Then $R \cdot C > 0$ for all curves not contained in the center of π . On the other hand, there are finitely many curves $B_1, \dots, B_N$ such that $R \cdot B_j \leq 0$. These “bad” curves have to be taken into account.

We are able to get around this difficulty since every splitting takes place in the fixed surface S_1 . Inside this surface we will not have any curves with “negative” homology.

We consider the normalisation

$$\eta : \tilde{S}_1 \rightarrow S_1.$$

We define a family $(\tilde{C}_t^{(k)})$ in $\tilde{S}_1$ by letting $\tilde{C}_t^{(k)}$ be the strict transform of $C_t^{(k)}$ in $\tilde{S}_1$ for general t and then take closure in the cycle space. Let $\tilde{C}^{(k)}$ be the strict transform of $C^{(k)}$ in $\tilde{S}_1$. Then we obtain a splitting

$$\tilde{C}_{t_k}^{(k)} = \tilde{C}^{(k+1)} + \tilde{R}_k$$

for some t_k . Inductively we find

$$\tilde{C}^{(m)} \equiv \tilde{C}_{t_m}^{(m)} = \tilde{C}^{(m+1)} + \tilde{R}_m,$$

for all $m \in \mathbb{N}$. Here $\equiv$ denotes homology equivalence in $\tilde{S}_1$. It follows that

$$\tilde{C}^{(1)} \equiv \tilde{C}^{(m)} + \sum_{j=1}^{m-1} \tilde{R}_j,$$

i.e., $\tilde{C}^{(1)}$ is homology equivalent to a sum of arbitrary many effective curves in $\tilde{S}_1$. This contradicts Lemma 3.4. $\square$

We now prove the two technical lemmata used in the proof of Proposition 3.3 above.

Lemma 3.4. *Let S be a compact connected normal Moishezon surface. Then there exists a linear map $\varphi : H_2(S, \mathbb{Q}) \rightarrow \mathbb{Q}$ such that $\varphi([C]) \geq 1$ for all classes of irreducible curves C in S .*

Proof. It suffices to construct φ on the subspace V of $H_2(S; \mathbb{Q})$ generated by classes of irreducible curves. Let $\sigma : \hat{S} \rightarrow S$ be a desingularisation of S and note that the surface $\hat{S}$ is projective. We let $\hat{H}$ be an ample divisor on $\hat{S}$ and $\sigma_*(\hat{H}) = H$ be its push-down to S . Using the intersection theory on normal surfaces established in [Mum61] and [Sak84], we define

$$\varphi([C]) = C \cdot H = (\sigma^*C) \cdot (\sigma^*H).$$

Here σ^*D denotes the sum $\overline{D} + \sum a_i E_i$ of the strict transform $\overline{D}$ of the divisor D in $\hat{S}$ and an appropriately weighted sum of the exceptional curves of σ .

In order to check that φ is well-defined on homology classes, it suffices to show that $c_1(\mathcal{O}(\sigma^*C)) = 0$ for every C with $[C] = 0 \in H_2(S, \mathbb{Q})$. Following the notation and results presented in [Sak84], Section 3, this is equivalent to $c_1(\mathcal{O}(C)) \in \ker(\sigma^*) = \ker(\eta_S) \subset H^2(S, \mathbb{Q})$. Here $\eta_S : H^2(S, \mathbb{Q}) \rightarrow H_2(S, \mathbb{Q})$ denotes the Poincaré homomorphism on S . We may write

$$\eta_S(c_1(\mathcal{O}(C))) = \sigma_*(\eta_{\hat{S}}(c_1(\widetilde{\mathcal{O}(C)})))$$

for the Poincaré isomorphism $\eta_{\hat{S}} : H^2(\hat{S}, \mathbb{Q}) \rightarrow H_2(\hat{S}, \mathbb{Q})$ on $\hat{S}$ and $\widetilde{\mathcal{O}(C)} = \mathcal{O}(\sigma^*C)$. Since $\eta_{\hat{S}}(c_1(\mathcal{O}(\sigma^*C))) = [\sigma^*C]$ on the smooth surface $\hat{S}$, we conclude $\eta_S(c_1(\mathcal{O}(C))) = \sigma_*[\sigma^*C] = [C] = 0$ and obtain the desired vanishing.

It remains to check that $\varphi([C]) \geq 1$ for all classes of irreducible curves C in S . We have

$$\varphi([C]) = (\sigma^*C) \cdot (\sigma^*H) = (\sigma^*C) \cdot (\overline{H} + \sum b_i E_i).$$

Since $(\sigma^*C) \cdot E_j = 0$ for all j by definition of σ^*C (cf. [Sak84, Section 1]), we conclude

$$\varphi([C]) = (\sigma^*C) \cdot \overline{H} = (\overline{C} + \sum a_i E_i) \cdot \overline{H}.$$

Since $\overline{H} = \hat{H}$ is ample and C is effective, in particular $\overline{C}$ is effective and $a_i > 0$ for all i , the desired inequality follows. $\square$

Lemma 3.5. *Let S be an irreducible Moishezon surface with a covering family $(C_t)_{t \in T}$ of (rational) curves. Suppose $\dim T \geq 4$. Let $T' \subset T$ be the subset of those t for which C_t splits. Then $\dim T' \geq 2$.*

Proof. Let $x \in S$ and $T(x) = \{t \in T \mid x \in C_t\}$. Since $\dim T \geq 4$ by assumption, we have $\dim T(x) \geq 3$. (Consider the graph $p : Z \rightarrow S$ of the family $(C_t)_{t \in T}$ and observe that $\dim(p^{-1}(x)) \geq 3$ and $q : Z \rightarrow T$ restricted to $p^{-1}(x)$ is finite.) The same dimension count substituting T by $T(x)$ shows that

$$\dim(T(x) \cap T(x')) \geq 2$$

for $x, x' \in S$. Hence there exists a 2-dimensional subfamily through x, x' and therefore we obtain a 1-dimensional subfamily through x and x' such that all members split. In other words

$$\dim(T' \cap T(x)) \geq 1.$$

Varying x we conclude $\dim T' \geq 2$. $\square$

Having established Proposition 3.3, it remains to show that the rational quotient $r : X \rightarrow T$ is actually a $\mathbb{P}_1$ -bundle.

Proposition 3.6. *Assume that the rational quotient $r : X \rightarrow T$ is holomorphic and equidimensional. Then r is a $\mathbb{P}_1$ -bundle, T is smooth and $X = \mathbb{P}(T_T)$.*

Proof. As a first step, we show that the fibers of r must be irreducible. Assume the contrary and let $r^{-1}(t_0) = l_{t_0} = C^{(1)} + R$ be a reducible fiber such that $L \cdot C^{(1)} \geq 1$. Then $C^{(1)}$ deforms in an at least 2-dimensional family, hence $C^{(1)}$ is a member of $(l_t)_{t \in T}$, i.e., $C^{(1)} = l_{t_1}$ for a suitable $t_1 \in T$. Since r is holomorphic, this is only possible when $t_0 = t_1$, a contradiction.

So $r : X \rightarrow T$ is a holomorphic, equidimensional map of normal complex spaces and every fiber of r is a reduced, irreducible rational curve. Now the arguments of [Kol96, Theorem II.2.8] can be adapted to our situation and it follows that r is a $\mathbb{P}_1$ -bundle. In particular, T is smooth and $X = \mathbb{P}(T_T)$. $\square$

Recall that a surface in class $\mathcal{C}$ is Kähler. In total we have shown:

Theorem 3.7. *Let X be a compact contact threefold in class $\mathcal{C}$. If the rational quotient has dimension 2, then X is Kähler and of the form $\mathbb{P}(T_Y)$ with a Kähler surface Y . The projection $X \rightarrow Y$ is the rational quotient, i.e., Y is not uniruled.*

4. THE CASE OF A 1-DIMENSIONAL RATIONAL QUOTIENT

In this section we assume that X is a compact contact threefold in class $\mathcal{C}$ with contact line bundle L and a rational quotient $r : X \dashrightarrow Q$ of dimension $\dim Q = 1$. Then necessarily X is Moishezon and Q is a smooth curve B of genus at least 1. We observe that $r : X \rightarrow B$ is holomorphic. Our aim is to show that X is of the form $X = \mathbb{P}(T_Y)$ for some surface Y . The surface Y is then necessarily Moishezon, and since a smooth Moishezon surface is projective, we are going to show directly that X is projective.

Let B_0 be the set of points b in B such that $r^{-1}(b) = X_b$ is smooth.

Lemma 4.1. *Let $b \in B_0$. Then X_b is a Hirzebruch surface $X_b = \mathbb{P}(\mathcal{O}_{\mathbb{P}_1} \oplus \mathcal{O}_{\mathbb{P}_1}(-e))$ with $e > 0$ even.*

Proof. By adjunction $K_{X_b} = -2L|_{X_b}$, hence X_b is minimal. Moreover, X_b cannot be a projective plane $\mathbb{P}_2$ or a Hirzebruch surface with odd e . To exclude the quadric ($e = 0$), observe that X_b is not F -integral, i.e., the restriction of the contact form θ to X_b does not vanish identically, and hence

$$\theta|_{X_b} \in H^0(X_b, \Omega_{X_b}^1 \otimes L|_{X_b}) \neq 0,$$

which is impossible for $X_b = \mathbb{P}_1 \times \mathbb{P}_1$. $\square$

Every smooth fiber X_b of r has a uniquely defined non-splitting 1-dimensional family of rational curves, namely the ruling lines. All these rational curves together give rise to a 2-dimensional family $(l_y)_{y \in Y}$ of rational curves in X , where Y is the irreducible component of the cycle space parametrising generically the ruling lines. We obtain an almost holomorphic map $\pi : X \dashrightarrow Y$ and a holomorphic map $g : Y \rightarrow B$, $g(y) = r(l_y)$,

$$\begin{array}{ccc} X & \dashrightarrow^\pi & Y \supset Y_0 \\ & & \downarrow g \\ & & B \supset B_0 \end{array}$$

such that $Y_0 = g^{-1}(B_0)$ is a $\mathbb{P}_1$ -bundle over B_0 and π is again a $\mathbb{P}_1$ -bundle over Y_0 . In fact, if Y_b is the fiber over $b \in B_0$, then $X_b = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(-e_b)|Y_b)$.

The technical key to the main result of this section is

Proposition 4.2. *The family $(l_y)_{y \in Y}$ does not split.*

Proof. Suppose (l_y) splits. Then there exists a point $y_0 \in Y$, an irreducible curve $C^{(1)}$ with $L \cdot C^{(1)} \geq 1$ and an effective curve R such that $l_{y_0} = C^{(1)} + R$. As seen before the curve $C^{(1)}$ deforms in an at least 2-dimensional family $(C_t^{(1)})_{t \in T_1}$.

(1) Let us first consider the case $L \cdot C_t^{(1)} = 1$. If the family $(C_t^{(1)})_{t \in T_1}$ covers a surface, we find a 1-dimensional subfamily through a fixed point and contradict Lemma 2.6. If $(C_t^{(1)})_{t \in T_1}$ covers all of X , we recover the original family (l_y) as follows: notice that the general $C_t^{(1)}$ will be an irreducible rational curve in a smooth fiber X_b and $-K_{X_b} \cdot C_t = 2$ by adjunction. This implies that the general curve $C_t^{(1)}$ must be a ruling line, i.e., the general curve C_t must be a curve l_t . This may now be excluded using Lemma 2.6 by the same arguments as in Proposition 3.3.

(2) Having ruled out $L \cdot C_t^{(1)} = 1$, we consider the case $L \cdot C_t^{(1)} \geq 2$, i.e.,

$$-K_X \cdot C_t^{(1)} = 2L \cdot C_t^{(1)} \geq 4.$$

(2a) Assume that the family $(C_t^{(1)})_{t \in T_1}$ covers all of X and choose a general point $x \in X$. Dimension count shows that there is a 2-dimensional subfamily $(C_t^{(1)})_{t \in T_1(x)}$ through the point x , necessarily filling a rational surface S , which must be a fiber of r . Since x is general, there is a $b \in B$ with X_b smooth such that $S = X_b$. The family $(C_t^{(1)})_{t \in T_1(x)}$ splits as $C_{t_1}^{(1)} = C^{(2)} + R_1$ with $L \cdot C^{(2)} \geq 1$. If $L \cdot C^{(2)} \geq 2$ we repeat to whole process. Assume that $L \cdot C^{(k)} \geq 2$ for all k . Then, we obtain a

decomposition of the homology class of $C_{t_1}^{(1)}$ as a sum of arbitrary many effective curves $C_{t_1}^{(1)} \equiv \sum_{k=1}^K C^{(k)} + R_{K-1}$. As we can always choose a subfamily through a point of S , we can assume that $\sum_{k=1}^K C^{(k)} + R_{K-1} \subset S$ for all K . Calculating the degree with respect to an ample line bundle H on S , we obtain a contradiction. Hence at some stage the procedure has to stop, i.e. $L \cdot C_t^{(k)} = 1$ and we conclude by Lemma 2.6.

(2b) It remains to consider the case where $L \cdot C_t^{(1)} \geq 2$ and the family $(C_t^{(1)})$ covers a surface S , which is a component of a fiber X_b of r . The family must split and we choose a splitting component $C^{(2)}$ such that $L \cdot C^{(2)} \geq 1$. If $L \cdot C^{(2)} = 1$, we are done again; if $L \cdot C^{(2)} \geq 2$, we obtain an at least 4-dimensional family $(C_t^{(2)})$. If this family covers X , we are done by the arguments of (2a) applied to $(C_t^{(2)})$, instead of $(C_t^{(1)})$. Otherwise, the family $(C_t^{(2)})$ fills a component S' of the same fiber X_b , and as in the proof of Proposition 3.3, by choosing $C^{(2)}$ carefully, we may assume that $S' = S$. Now we are in completely the same situation as in the proof of Proposition 3.3 and proceed as described there. $\square$

In order to apply Proposition 4.2, we consider the normalised graph $p : Z \rightarrow X$ of the family $(l_y)_{y \in Y}$.

$$\begin{array}{ccc} Z & \xrightarrow{p} & X \\ & \searrow q & \swarrow \pi \\ & & Y \end{array}$$

Proposition 4.3. *The map $p : Z \rightarrow X$ is biholomorphic.*

Proof. The map p is generically biholomorphic, hence by Zariski's main theorem, it suffices to show that p does not have positive-dimensional fibers. So suppose that $\dim p^{-1}(x) = 1$. Then there exists a 1-dimensional subfamily $(l_y)_{y \in Y(x)}$ through x with all l_y irreducible by the previous proposition. Since $L \cdot l_y = 1$, we contradict Lemma 2.6. $\square$

As before in Proposition 3.6 we conclude:

Corollary 4.4. *The map $\pi : X \rightarrow Y$ is a $\mathbb{P}_1$ -bundle.*

We may now apply Lemma 4.6 below to the map $\pi : X \rightarrow Y$ and have shown:

Theorem 4.5. *Let X be a compact contact threefold in class $\mathcal{C}$ with 1-dimensional rational quotient B . Then X is projective and there is a smooth projective surface Y with a $\mathbb{P}_1$ -fibration $X \rightarrow Y$ such that $X \simeq \mathbb{P}(T_Y)$.*

Lemma 4.6. *Let Y be a complex manifold of dimension $n + 1$ and $\pi : X \rightarrow Y$ be a $\mathbb{P}_n$ -bundle. If X is a contact manifold, then $X \simeq \mathbb{P}(T_Y)$.*

Proof. Let $Z \simeq \mathbb{P}_n$ be a fiber of π . Adjunction implies that the contact line bundle L restricted to Z fulfils $L|_Z = \mathcal{O}_Z(1)$. Setting $\mathcal{E} = \pi_*(L)$, we conclude

$$(X, L) \simeq (\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)).$$

Now the last part of the proof of Theorem 2.12 in [KPSW00] can be applied and shows that $\mathcal{E} \simeq T_Y$. $\square$

REFERENCES

- [BDPP04] Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell, *The pseudo-effective cone of a compact Kähler manifold of negative Kodaira dimension*, arXiv:math/0405285, 2004.
- [Bea98] Arnaud Beauville, *Fano contact manifolds and nilpotent orbits*, Comment. Math. Helv. **73** (1998), no. 4, 566–583.
- [Bea00] ———, *Complex manifolds with split tangent bundle*, Complex analysis and algebraic geometry, volume in memory of Michael Schneider, eds. T. Peternell and F.O. Schreyer, de Gruyter, Berlin, 2000, pp. 61–70.
- [Bru06] Marco Brunella, *A positivity property for foliations on compact Kähler manifolds*, Internat. J. Math. **17** (2006), no. 1, 35–43.
- [Cam80] Frederic Campana, *Algébricité et compacité dans l'espace des cycles d'un espace analytique complexe*, Math. Ann. **251** (1980), no. 1, 7–18.
- [Deb01] Olivier Debarre, *Higher-dimensional algebraic geometry*, Universitext, Springer-Verlag, New York, 2001.
- [Dem02] Jean-Pierre Demailly, *On the Frobenius integrability of certain holomorphic p -forms*, Complex geometry (Göttingen, 2000), volume in honour of H. Grauert, eds. I. Bauer et al., Springer, Berlin, 2002, pp. 93–98.
- [GHS03] Tom Graber, Joe Harris, and Jason Starr, *Families of rationally connected varieties*, J. Amer. Math. Soc. **16** (2003), no. 1, 57–67.
- [Keb01] Stefan Kebekus, *Lines on contact manifolds*, J. Reine Angew. Math. **539** (2001), 167–177.
- [Kol96] János Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 32, Springer-Verlag, Berlin, 1996.
- [KPSW00] Stefan Kebekus, Thomas Peternell, Andrew J. Sommese, and Jarosław A. Wiśniewski, *Projective contact manifolds*, Invent. Math. **142** (2000), no. 1, 1–15.
- [LeB95] Claude LeBrun, *Fano manifolds, contact structures, and quaternionic geometry*, Internat. J. Math. **6** (1995), no. 3, 419–437.
- [MM86] Yoichi Miyaoka and Shigefumi Mori, *A numerical criterion for uniruledness*, Ann. of Math. (2) **124** (1986), no. 1, 65–69.
- [Mum61] David Mumford, *The topology of normal singularities of an algebraic surface and a criterion for simplicity*, Inst. Hautes Études Sci. Publ. Math. (1961), no. 9, 5–22.
- [Pet01] Thomas Peternell, *Towards a Mori theory on compact Kähler threefolds. III*, Bull. Soc. Math. France **129** (2001), no. 3, 339–356.
- [Sak84] Fumio Sakai, *Weil divisors on normal surfaces*, Duke Math. J. **51** (1984), no. 4, 877–887.

MATHEMATISCHES INSTITUT, UNIVERSITÄT BAYREUTH, 95440 BAYREUTH, GERMANY
 KRISTINA.FRANTZEN@UNI-BAYREUTH.DE, THOMAS.PETERNELL@UNI-BAYREUTH.DE